

Gene Section

Review

DNMT3B (DNA (cytosine-5)-methyltransferase 3 beta)

Riadh Ben Gacem, Olfa Ben Abdelkrim, Sonia Ziadi, Mounir Trimeche

Department of Pathology, Farhat-Hached Hospital, Sousse, Tunisia (RB, OB, SZ, MT)

Published in Atlas Database: July 2012

Online updated version: <http://AtlasGeneticsOncology.org/Genes/DNMT3BID40350ch20q11.html>

DOI: 10.4267/2042/48489

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.

© 2013 Atlas of Genetics and Cytogenetics in Oncology and Haematology

Identity

Other names: ICF, ICF1, M.HsaIIIB

HGNC (Hugo): DNMT3B

Location: 20q11.21

Note

DNA (cytosine-5)-methyltransferase 3B (DNMT3B) is one of the three known DNA methyltransferases with catalytic activity.

This enzyme is responsible for the de novo methylation of DNA, particularly during embryogenesis (Okano et al., 1999). In fact, DNMT3B adds methyl groups to CpG dinucleotides of unmethylated DNA for establishment of new methylation pattern in genomic DNA. DNMT3B level is profoundly increased in various tumor cell lines and in numerous types of human cancers,

indicating that it plays an important role in tumorigenesis (Turek-Plewa and Jagodzinski, 2005).

DNA/RNA

Description

DNMT3B gene is located on the long arm of chromosome 20 at position 11.2 and is composed of 23 exons and 22 introns. This gene encodes for a protein of 853 aa. DNMT3B gene is abundantly expressed in embryonic stem cells, but its expression is decreased upon their differentiation and remains low in adult somatic tissues.

Transcription

Primary transcript of DNMT3B gene can be spliced five different mRNA isoforms that are DNMT3B1, DNMT3B2, DNMT3B3, DNMT3B4 and DNMT3B5.




Figure 1. DNMT3B gene is located on the long arm of chromosome 20 at position 11.2 from the base pair 31350190 to base pair 31397161. It is composed of 23 exons and 22 introns.




Figure 2. A: The general structure of DNMT3B protein. The N-terminal domain contains a proliferating cell nuclear antigen-binding domain, a nuclear localization signal, a tetrapeptide PWWP, essential for DNMT binding to chromatin, an ATRX cysteine-rich zinc finger DNA-binding motif and a polybromo homology domain (PHD) targeting DNMT3B to the replication foci. The C-terminal catalytic domain of DNMT3B is characterized by the presence of 6 conserved amino acid motifs, namely I, IV, VI, VIII, IX and X. Motifs I and X form S-adenosylmethionine binding site, motif IV binds cytosine at the active site, motif VI possesses glutamyl residue donating protons, and motif IX maintains the structure of the target recognition domain (TRD) usually located between motifs VIII and IX, that make base-specific contacts in the major groove of DNA (Turek-Plewa and Jagodzinski, 2005). **B:** The results from alternative splicing of exons 10, 21 and/or 22, have reported several different transcripts of DNMT3B. These isoforms are: DNMT3B1, DNMT3B2, DNMT3B3, DNMT3B4 and DNMT3B5. DNMT3B1 and DNMT3B2, contain all the highly conserved motifs I, IV, VI, IX and X, are enzymatically active in DNA methyltransferase assays. DNMT3B3, which lacks parts of motif IX, appears to be inactive enzyme. DNMT3B4 and DNMT3B5, encode truncated proteins that lack motifs IX and X, are presumably inactive as well. The lack motifs in DNMT3B3, DNMT3B4 and DNMT3B5 are a consequence of lack 21-22, 21 or 22 exons, respectively (Xie et al., 1999; Okano et al., 1999).

Protein

Note

DNMT3B protein is structured with N-terminal regulatory and C-terminal catalytic domains that are linked by repeated GK dipeptides (Glycine-Lysine-repeats). The N-terminal domain possesses nuclear localization signal sequence (NLS) responsible for nucleus DNMT3B localization. The N-terminal domain plays a regulatory role, and contains a proliferating cell nuclear antigen-binding domain. This N-domain also contains a cysteine rich zinc finger DNA binding motif (ATRX), a polybromo homology domain (PHD) targeting DNMT3B to the replication foci and a PWWP tetrapeptide

chromatin-binding domain (Margot et al., 2000). The two halves are packed against each other to form a single structural module that exhibits a prominent positive electrostatic potential. The PWWP domain of DNMT3B alone binds DNA in vitro, probably through its basic surface, and DNMT3B binds DNA stronger than a mutant without the domain. In addition, the PWWP domain seems to target the de novo methyltransferases to chromatin. The C-terminal domain contains six conservative motifs I, IV, VI, VIII, IX and X. Motif I and X are involved in the formation of the S-adenosylmethionine binding site. Motif IV binds the substrate cytosine at the active site. Motif VI contains the glutamyl residue serving as a proton

donor. Motif IX maintains the structure of the target recognition domain (TRD). Motif VIII's function is unclear (Xie et al., 1999).

The intracellular distribution of DNMT3B enzyme is rather dynamic throughout the cell cycle, indeed this enzyme is diffusely distributed throughout the nucleoplasm during most of G1, associates with subnuclear sites of DNA replication during S-phase (Leonhardt et al., 1992), and binds to chromatin, with preference to pericentric heterochromatin, during G2 and M-phases (Easwaran et al., 2005).

Description

853 amino acids; 95751 Da.

Localisation

Nucleus.

Function

DNMT3B is expressed at very low levels in most tissues except the testis, thyroid and bone marrow (Xie et al., 1999). DNMT3B level is profoundly increased in various tumor cell lines, indicating that it plays an important role in tumorigenesis (Robertson et al., 1999; Hermann et al., 2004). DNA methylation is a covalent chemical modification, resulting in the addition of a methyl (CH₃) group at the carbon 5 position of the cytosine ring. DNMT3B uses S-adenosyl-L-methionine (AdoMet) as the source of the methyl group being transferred to the DNA bases. The methyl group of S-adenosylomethionine is bound to a sulphonium atom, which thermodynamically destabilizes the molecule and makes the relatively inert methylthiol of the methionine moiety very reactive towards nucleophilic attack by nitrogen, oxygen and sulphur atoms or activated C atoms (carbanions).

DNMT3B may also interact with DNMT1 and activate HDAC1, which deacetylates histones and represses gene transcription.

This indicates that DNMT3B may be involved in chromatin remodeling associated with the modulation of gene transcription. DNMT3B can also effectively methylate C to m⁵C post-replicatively in unmethylated DNA. During or after replication, DNA regions may bind sequence-specific proteins which block the attachment of the methyl group to CpG dinucleotide and the formation of methylation patterns unique for each tissue.

The existence of DNMT3B isoforms suggests that other factors can be involved in the binding of DNMT3B to a particular DNA region. Consequences of alternative splicing on DNMT3B's ability to interact with DNMT3L. DNMT3L stimulates the catalytic activity of DNMT3A and DNMT3B methyltransferases (Suetake et al., 2004). DNMT3L binds to the carboxyl-terminal part of DNMT3B and increases the level of activity of this enzyme (Suetake et al., 2004).

Homology

DNMT3B exhibit a high degree of primary structure homology with DNMT3A. Structurally, DNMT3B and DNMT3A have a similar organization, with conserved domains: the long N-terminal region contains a PWWP domain and a cysteine-rich PHD zinc finger domain and the C-terminal catalytic domain. The same NLS and ATRX sequences were also founded in the DNMT3B and DNMT3A enzymes.

Mutations

Note

ICF syndrome (standing for Immunodeficiency, Centromere instability and Facial anomalies syndrome) is a rare autosomal recessive immune disorder caused by mutations of the gene DNMT3B. Therefore, the DNMT3B mutations, like A766P and R840Q, observed in patients are the likely underlying cause of their ICF phenotypes (Xie et al., 2006).

This hereditary syndrome is characterized by centromeric instability of chromosomes 1, 9, and 16 is associated with abnormal hypomethylation of CpG sites in their pericentromeric satellite regions (Hansen et al., 1999). At the molecular level, in patient DNA, sequences such as the pericentromeric classic satellite repeats of pericentromeric regions are hypomethylated, which attributes to reduced enzymatic activity of the mutant proteins (Jeanpierre et al., 1993; Gowher and Jeltsch, 2002).

Several polymorphisms were detected at the promoter region of DNMT3B gene and may influence its activity in DNA methylation and increase the susceptibility to several cancers. DNMT3B promoter polymorphisms influence on promoter hypermethylation of genes in other interacting pathways, such as cell cycle, apoptosis, or other DNA repair pathways and may increase level of DNA damage which contributes to an increased risk for cancers. For example, C>T (rs406193), -283T>C (rs121909506), -579G>T (rs2424909), -579G>T (rs2235758) and G39179T DNMT3B polymorphisms influence DNMT3B expression, thus contributing to the genetic susceptibility to different cancers (Lee et al., 2005; Fan et al., 2008; Montgomery et al., 2004; de Vogel et al., 2009; Daraei et al., 2011).

The common DNMT3B -149C>T (rs2424913) polymorphism was found to significantly increase DNMT3B promoter activity. This up-regulates the expression of the gene and may in turn lead to aberrant methylation of CpG islands in some tumor suppressor genes (Montgomery et al., 2004). It was associated with an increased risk for lung cancer (Shen et al., 2002), prostate cancer (Singal et al., 2005), and colorectal polyps, including colorectal adenomas (Jung et al., 2008), or with prognosis of head and neck cancer (Wang et al., 2004).




Figure 3. DNMT3B overexpression in breast cancer cells as detected by immunocytochemistry (original magnification x400).

Implicated in

Lung cancer

Note

DNMT3B may play an oncogenic role during tumorigenesis, and its genetic variants have been consistently associated with risk of several cancers. Overexpression of DNMT3B may result in promoter hypermethylation of multiple tumor suppressor genes and ultimately lead to lung tumorigenesis and poor prognosis especially among smoking patients (Lin et al., 2007; Palakurthy et al., 2009).

Polymorphisms and haplotypes of the DNMT3B gene may influence DNMT3B activity on DNA methylation, thereby modulating the susceptibility to lung cancer. The -283T>C and -579G>T polymorphisms in the DNMT3B promoter, and their haplotypes were significantly associated with the risk of adenocarcinoma of the lung (Lee et al., 2005). These mutations up-regulate DNMT3B expression, resulting in a predisposition towards aberrant de novo methylation of CpG islands in tumor suppressor genes and DNA repair genes (Lee et al., 2005; Shen et al., 2002).

Breast cancer

Note

DNMT3B expression levels are significantly higher

in breast cancers than in normal breast tissues, suggesting a role of this enzyme in breast tumorigenesis (Fig. 3) (Girault et al., 2003; Butcher and Rodenhiser, 2007; Ben Gacem et al., 2012). DNMT3B overexpression is correlated to the epigenetic inactivation of several tumor suppressor genes and to the aggressive phenotype in breast tumors (Roll et al., 2008; Ben Gacem et al., 2012). Some polymorphisms are associated to the development of breast tumors. DNMT3B C46359T polymorphism has been correlated to women with early-onset breast cancer, bilateral breast cancer and with a family history of the disease (Montgomery et al., 2004; Wang et al., 2004).

Leukemia and lymphomas

Note

DNMT3B is substantially overexpressed in leukemia cells than in normal bone marrow cells (Mizuno et al., 2001). DNMT3B overexpression was correlated to hypermethylation of several tumor related genes in diffuse large B-cell lymphomas and in B-cell chronic lymphocytic leukemia (Amara et al., 2010; Kn et al., 2004). In B-cell chronic lymphocytic leukemia, DNMT3B overexpression was identified as an independent prognostic factor for predicting shortened survival of patients (Amara et al., 2010).

Hepatocellular carcinoma

Note

DNA hypomethylation on pericentromeric satellite regions is one of the earliest events during hepatocarcinogenesis. Overexpression of DNMT3B4 isoform and elevation of the ratio of DNMT3B4 mRNA to DNMT3B3 mRNA were both significantly correlated with the degree of DNA hypomethylation on pericentromeric satellite regions in hepatocellular carcinoma (Saito et al., 2002). DNMT3B4 overexpression induces DNA demethylation on pericentromeric satellite regions even in precancerous stages and may play critical roles in the development of hepatocellular carcinoma through chromosomal instability and aberrant expression of cancer-related genes (Saito et al., 2002).

Colorectal cancer

Note

The levels of DNMTs mRNA/protein in colorectal carcinomas are significantly higher than in noncancerous colorectal tissues and the highest expression range was observed with DNMT3B (Eads et al., 1999; Linhart et al., 2007; Noshio et al., 2009; Huidobro et al., 2012). The CpG island methylator phenotype (CIMP) in colorectal cancer is characterized by a widespread CpG island methylation. DNMT3B contributes to CpG island methylation, which may eventually lead to the development of CIMP-high colorectal cancer (Noshio et al., 2009; Schmidt et al., 2007).

In hereditary nonpolyposis colorectal cancer, the DNMT3B gene contains a C-to-T single nucleotide polymorphism -149 bp from the transcriptional start site that may result in increased promoter activity of the gene (Shen et al., 2002). The DNA mismatch repair mutation carriers who also carried at least one T allele for the DNMT3B promoter region C-to-T polymorphism had a 2-fold higher risk for colorectal cancer by year than those who were homozygous for the wild-type DNMT3B allele (Jones et al., 2006).

Facial anomalies syndrome (ICF)

Note

Facial anomalies syndrome (ICF) is an autosomal recessive disease. DNMT3B gene is often the site of ICF mutations (Ehrlich et al., 2006). ICF syndrome presents with variable combined immunodeficiency, mild facial anomalies and extravagant cytogenetic abnormalities with largely affect the pericentric regions of chromosomes 1, 9 and 16. These pericentric regions contain a type of satellite DNA termed classical satellite, or satellites 2 and 3. It is normally heavily methylated, but is nearly completely unmethylated in DNA of ICF patients. In addition to classical satellite, demethylation of DNA in ICF patients is also seen at CpG islands on the inactivate X chromosome in females (Ehrlich et al., 2008).

References

- Leonhardt H, Page AW, Weier HU, Bestor TH. A targeting sequence directs DNA methyltransferase to sites of DNA replication in mammalian nuclei. *Cell*. 1992 Nov 27;71(5):865-73
- Jeanpierre M, Turleau C, Aurias A, Prieur M, Ledest F, Fischer A, Viegas-Pequignot E. An embryonic-like methylation pattern of classical satellite DNA is observed in ICF syndrome. *Hum Mol Genet*. 1993 Jun;2(6):731-5
- Eads CA, Danenberg KD, Kawakami K, Saltz LB, Danenberg PV, Laird PW. CpG island hypermethylation in human colorectal tumors is not associated with DNA methyltransferase overexpression. *Cancer Res*. 1999 May 15;59(10):2302-6
- Hansen RS, Wijmenga C, Luo P, Stanek AM, Canfield TK, Weemaes CM, Gartner SM. The DNMT3B DNA methyltransferase gene is mutated in the ICF immunodeficiency syndrome. *Proc Natl Acad Sci U S A*. 1999 Dec 7;96(25):14412-7
- Okano M, Bell DW, Haber DA, Li E. DNA methyltransferases Dnmt3a and Dnmt3b are essential for de novo methylation and mammalian development. *Cell*. 1999 Oct 29;99(3):247-57
- Robertson KD, Uzvolgyi E, Liang G, Talmadge C, Sumegi J, Gonzales FA, Jones PA. The human DNA methyltransferases (DNMTs) 1, 3a and 3b: coordinate mRNA expression in normal tissues and overexpression in tumors. *Nucleic Acids Res*. 1999 Jun 1;27(11):2291-8
- Xie S, Wang Z, Okano M, Nogami M, Li Y, He WW, Okumura K, Li E. Cloning, expression and chromosome locations of the human DNMT3 gene family. *Gene*. 1999 Aug 5;236(1):87-95
- Margot JB, Aguirre-Arteta AM, Di Giacco BV, Pradhan S, Roberts RJ, Cardoso MC, Leonhardt H. Structure and function of the mouse DNA methyltransferase gene: Dnmt1 shows a tripartite structure. *J Mol Biol*. 2000 Mar 24;297(2):293-300
- Mizuno S, Chijiwa T, Okamura T, Akashi K, Fukumaki Y, Niho Y, Sasaki H. Expression of DNA methyltransferases DNMT1, 3A, and 3B in normal hematopoiesis and in acute and chronic myelogenous leukemia. *Blood*. 2001 Mar 1;97(5):1172-9
- Gowher H, Jeltsch A. Molecular enzymology of the catalytic domains of the Dnmt3a and Dnmt3b DNA methyltransferases. *J Biol Chem*. 2002 Jun 7;277(23):20409-14
- Saito Y, Kanai Y, Sakamoto M, Saito H, Ishii H, Hirohashi S. Overexpression of a splice variant of DNA methyltransferase 3b, DNMT3b4, associated with DNA hypomethylation on pericentromeric satellite regions during human hepatocarcinogenesis. *Proc Natl Acad Sci U S A*. 2002 Jul 23;99(15):10060-5
- Shen H, Wang L, Spitz MR, Hong WK, Mao L, Wei Q. A novel polymorphism in human cytosine DNA-methyltransferase-3B promoter is associated with an increased risk of lung cancer. *Cancer Res*. 2002 Sep 1;62(17):4992-5
- Girault I, Tozlu S, Lidereau R, Bièche I. Expression analysis of DNA methyltransferases 1, 3A, and 3B in sporadic breast carcinomas. *Clin Cancer Res*. 2003 Oct 1;9(12):4415-22
- Hermann A, Gowher H, Jeltsch A. Biochemistry and biology of mammalian DNA methyltransferases. *Cell Mol Life Sci*. 2004 Oct;61(19-20):2571-87

- Kn H, Bassal S, Tikellis C, El-Osta A. Expression analysis of the epigenetic methyltransferases and methyl-CpG binding protein families in the normal B-cell and B-cell chronic lymphocytic leukemia (CLL). *Cancer Biol Ther.* 2004 Oct;3(10):989-94
- Montgomery KG, Liu MC, Eccles DM, Campbell IG. The DNMT3B C-->T promoter polymorphism and risk of breast cancer in a British population: a case-control study. *Breast Cancer Res.* 2004;6(4):R390-4
- Suetake I, Shinozaki F, Miyagawa J, Takeshima H, Tajima S. DNMT3L stimulates the DNA methylation activity of Dnmt3a and Dnmt3b through a direct interaction. *J Biol Chem.* 2004 Jun 25;279(26):27816-23
- Wang L, Rodriguez M, Kim ES, Xu Y, Bekele N, El-Naggar AK, Hong WK, Mao L, Oh YW. A novel C/T polymorphism in the core promoter of human de novo cytosine DNA methyltransferase 3B6 is associated with prognosis in head and neck cancer. *Int J Oncol.* 2004 Oct;25(4):993-9
- Easwaran HP, Leonhardt H, Cardoso MC. Cell cycle markers for live cell analyses. *Cell Cycle.* 2005 Mar;4(3):453-5
- Lee SJ, Jeon HS, Jang JS, Park SH, Lee GY, Lee BH, Kim CH, Kang YM, Lee WK, Kam S, Park RW, Kim IS, Cho YL, Jung TH, Park JY. DNMT3B polymorphisms and risk of primary lung cancer. *Carcinogenesis.* 2005 Feb;26(2):403-9
- Singal R, Das PM, Manoharan M, Reis IM, Schlesselman JJ. Polymorphisms in the DNA methyltransferase 3b gene and prostate cancer risk. *Oncol Rep.* 2005 Aug;14(2):569-73
- Turek-Plewa J, Jagodziński PP. The role of mammalian DNA methyltransferases in the regulation of gene expression. *Cell Mol Biol Lett.* 2005;10(4):631-47
- Ehrlich M, Jackson K, Weemaes C. Immunodeficiency, centromeric region instability, facial anomalies syndrome (ICF). *Orphanet J Rare Dis.* 2006 Mar 1;1:2
- Jones JS, Amos CI, Pande M, Gu X, Chen J, Campos IM, Wei Q, Rodriguez-Bigas M, Lynch PM, Frazier ML. DNMT3b polymorphism and hereditary nonpolyposis colorectal cancer age of onset. *Cancer Epidemiol Biomarkers Prev.* 2006 May;15(5):886-91
- Xie ZH, Huang YN, Chen ZX, Riggs AD, Ding JP, Gowher H, Jeltsch A, Sasaki H, Hata K, Xu GL. Mutations in DNA methyltransferase DNMT3B in ICF syndrome affect its regulation by DNMT3L. *Hum Mol Genet.* 2006 May 1;15(9):1375-85
- Butcher DT, Rodenhiser DI. Epigenetic inactivation of BRCA1 is associated with aberrant expression of CTCF and DNA methyltransferase (DNMT3B) in some sporadic breast tumours. *Eur J Cancer.* 2007 Jan;43(1):210-9
- Lin RK, Hsu HS, Chang JW, Chen CY, Chen JT, Wang YC. Alteration of DNA methyltransferases contributes to 5'CpG methylation and poor prognosis in lung cancer. *Lung Cancer.* 2007 Feb;55(2):205-13
- Linhart HG, Lin H, Yamada Y, Moran E, Steine EJ, Gokhale S, Lo G, Cantu E, Ehrlich M, He T, Meissner A, Jaenisch R. Dnmt3b promotes tumorigenesis in vivo by gene-specific de novo methylation and transcriptional silencing. *Genes Dev.* 2007 Dec 1;21(23):3110-22
- Schmidt WM, Sedivy R, Forstner B, Steger GG, Zöchbauer-Müller S, Mader RM. Progressive up-regulation of genes encoding DNA methyltransferases in the colorectal adenoma-carcinoma sequence. *Mol Carcinog.* 2007 Sep;46(9):766-72
- Ehrlich M, Sanchez C, Shao C, Nishiyama R, Kehrl J, Kuick R, Kubota T, Hanash SM. ICF, an immunodeficiency syndrome: DNA methyltransferase 3B involvement, chromosome anomalies, and gene dysregulation. *Autoimmunity.* 2008 May;41(4):253-71
- Fan H, Zhang F, Hu J, Liu D, Zhao Z. Promoter polymorphisms of DNMT3B and the risk of colorectal cancer in Chinese: a case-control study. *J Exp Clin Cancer Res.* 2008 Jul 28;27:24
- Jung AY, Poole EM, Bigler J, Whitton J, Potter JD, Ulrich CM. DNA methyltransferase and alcohol dehydrogenase: gene-nutrient interactions in relation to risk of colorectal polyps. *Cancer Epidemiol Biomarkers Prev.* 2008 Feb;17(2):330-8
- Roll JD, Rivenbark AG, Jones WD, Coleman WB. DNMT3b overexpression contributes to a hypermethylator phenotype in human breast cancer cell lines. *Mol Cancer.* 2008 Jan 25;7:15
- de Vogel S, Wouters KA, Gottschalk RW, van Schooten FJ, de Goeij AF, de Bruïne AP, Goldbohm RA, van den Brandt PA, Weijnenberg MP, van Engeland M. Genetic variants of methyl metabolizing enzymes and epigenetic regulators: associations with promoter CpG island hypermethylation in colorectal cancer. *Cancer Epidemiol Biomarkers Prev.* 2009 Nov;18(11):3086-96
- Nosho K, Shima K, Irahara N, Kure S, Baba Y, Kirkner GJ, Chen L, Gokhale S, Hazra A, Spiegelman D, Giovannucci EL, Jaenisch R, Fuchs CS, Ogino S. DNMT3B expression might contribute to CpG island methylator phenotype in colorectal cancer. *Clin Cancer Res.* 2009 Jun 1;15(11):3663-71
- Palakurthy RK, Wajapeyee N, Santra MK, Gazin C, Lin L, Gobeil S, Green MR. Epigenetic silencing of the RASSF1A tumor suppressor gene through HOXB3-mediated induction of DNMT3B expression. *Mol Cell.* 2009 Oct 23;36(2):219-30
- Amara K, Ziadi S, Hachana M, Soltani N, Korbi S, Trimeche M. DNA methyltransferase DNMT3b protein overexpression as a prognostic factor in patients with diffuse large B-cell lymphomas. *Cancer Sci.* 2010 Jul;101(7):1722-30
- Daraei A, Salehi R, Mohamashem F. DNA-methyltransferase 3B 39179 G > T polymorphism and risk of sporadic colorectal cancer in a subset of Iranian population. *J Res Med Sci.* 2011 Jun;16(6):807-13
- Ben Gacem R, Hachana M, Ziadi S, Ben Abdelkarim S, Hidar S, Trimeche M. Clinicopathologic significance of DNA methyltransferase 1, 3a, and 3b overexpression in Tunisian breast cancers. *Hum Pathol.* 2012 Oct;43(10):1731-8
- Huidobro C, Urdinguio RG, Rodríguez RM, Mangas C, Calvanese V, Martínez-Camblor P, Ferrero C, Parra-Blanco A, Rodrigo L, Obaya AJ, Suárez-Fernández L, Astudillo A, Hernando H, Ballestar E, Fernández AF, Fraga MF. A DNA methylation signature associated with aberrant promoter DNA hypermethylation of DNMT3B in human colorectal cancer. *Eur J Cancer.* 2012 Sep;48(14):2270-81

This article should be referenced as such:

Ben Gacem R, Ben Abdelkrim O, Ziadi S, Trimeche M. DNMT3B (DNA (cytosine-5)-methyltransferase 3 beta). *Atlas Genet Cytogenet Oncol Haematol.* 2013; 17(2):88-93.
